

Timed Up & Go Test (TUG)

Research Report

Author: Anne Shumway-Cook, Sandy Brauer, and Marjorie Woollacott

Description of the Instrument

- Patients are timed (in seconds) when performing the TUG—3 conditions
 1. TUG alone—from sitting in a chair, stand up, walk 3 meters, turn around, walk back, and sit down..
 2. TUG Cognitive—complete the task while counting backwards from a randomly selected number between 20 and 100.
 3. TUG manual—complete the task while carrying a full cup of water.
- The time taken to complete the task is strongly correlated to level of functional mobility, (i.e. the more time taken, the more dependent in activities of daily living).
- **The cutoff levels for TUG is 13.5 seconds or longer with an overall correct prediction rate of 90%**; for TUG manual is 14.5 seconds or longer with a 90% correct prediction rate; and Tug cognitive is 15 seconds or longer with an overall correct prediction rate of 87%.

Form of instrument:

- Hazard/Risk Assessment Tools
- To identify/screen elderly individuals who are prone to falls
- Interrater reliability was very high, with $r=.98$, $.99$, and $.99$ for the TUG, TUGmanual, and TUGcognitive respectively
- The TUG alone correctly classified 13/15 fallers (87% sensitivity) and 13/15 nonfallers (87% specificity).

Validity Measures

- **Older adults who take longer than 14 seconds to complete the TUG have a high risk for falls. This cutoff is different from Podsiadlo and Richardson, which is 30 seconds.**

Reference

- Lundlin-Olsson, L., Nyberg, L., & Gustafson, Y. (1998). Attention, frailty, and falls: the effect of a manual task on basic mobility. *Journal of the American Geriatrics Society*, 46, 758-761.
- Podsiadlo, D., & Richardson, S. (1991). The timed “up & go”: A test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society*, 39, 142-148.
- Shumway-Cook, A., Brauer, S., & Woollacott, M. (2000). Predicting the probability for falls in community-dwelling older adults using the timed up & go test. *Physical Therapy*, 80(9), 896-903.